# 2014-2015 Influenza Report

### Week 14

April 5, 2015 - April 11, 2015


### About our flu activity reporting

MSDH relies upon selected sentinel health practitioners across the state to report the percentage of total patient visits consistent with an influenza-like illness (ILI: fever of 100°F or higher AND cough and/or sore throat). Also, providers are supplied with specimen collection kits. Samples are submitted to the Mississippi Public Health Laboratory for influenza PCR testing. Reports are used to estimate the state's ILI rate and the magnitude of the state's influenza activity. Reports represent only the distribution of flu in the state, not an actual count of all flu cases statewide. *Information is provisional only and may change depending on additional reporting from sentinel providers.* 

#### Content

- State ILI Surveillance
- Flu Testing Reports
- National and Mississippi Pediatric Mortality Surveillance
- **National ILI Surveillance**
- Appendix
  - Figure 1 (Statewide ILI Data, Current Season compared to Previous Season)
  - Figure 2 (Percentage of ILI Cases by Age Group, Mississippi)
  - Figure 3 (Mississippi ILI Rates, 2014-2015 and Previous Seasons)
  - Figure 4 (State ILI/URI Rates 2011-2015 (YTD))
  - Figure 5 (Comparison of Statewide ILI Rate to Positive Influenza Isolates by Type and Subtype, Mississippi)
  - Figure 6 (Number of Influenza-Associated Pediatric Deaths by Week of Death and Influenza Type, Nationwide)
  - <u>Figure 7</u> (Percentage of Influenza-Associated Pediatric Deaths by Influenza Type, Nationwide)
  - Figure 8 (Comparison of Nationwide ILI Rates to Mississippi ILI Rates)
  - o Figure 9 (CDC FluView Weekly Influenza Activity Map)

#### **State ILI Surveillance**



During week 14 (04/05/15-04/11/15), the overall state ILI rate (3.1%) was comparable to the previous week (3.2%), but was higher than this time last year (2.3%).

| Figure 1


Total number of patients treated by sentinel providers in the last three weeks. | Table 1

2014-2015 Influenza Season						
CDC Week	Week Ending	Number of ILI Reports	Total patients	ILI symptoms	ILI Rate (%)	
14	Apr. 11	137	18473	574	3.1%	
13	Apr. 04	152	18513	600	3.2%	
12	Mar. 28	149	18941	754	4.0%	

During week **14**, **two** districts (3 and 7) had an increase in ILI activity, while **two** districts (2 and 6) had a decrease. **Five** districts (1, 4, 5, 8, and 9) remained about the same. *Information is provisional only and may change depending on additional reporting from sentinel providers*. | **Table 2** 



MSDH District ILI Rates (%) 2014-2015					
District	Week 13	Week 14			
State	<i>3.2</i>	3.1			
I	1.4	1.5			
II	4.9	1.4			
III	4.6	7.1			
IV	2.9	3.3			
V	2.9	2.9			
VI	9.3	7.0			
VII	1.9	3.1			
VIII	1.7	1.3			
IX	4.3	4.2			




Since week
35, the
percentage
of reported
ILI cases has
been highest
among those
in the 5-24
years of age
group.
Overall, this


trend

continued, but during week **14**, the 0-4 years of age group and 5-24 years of age group had the same rate. | Figure 2

The 2014-15 state ILI rate was **above** the national and Region 4 baselines, but was **below** the state baseline, during week **14**. | Figure 3




## Mississippi ILI Rates 2011-2015 | Figure 4



#### **Flu Testing Reports**


From week **40** (week ending October 4<sup>th</sup>) through week **14** (week ending April 11<sup>th</sup>), 193 positive influenza samples were identified by MSDH. One hundred sixty-one (83%) samples were identified as influenza A (H3), eight (4%) as influenza A (H1), and twenty-four (12%) were identified as influenza B. The last positive influenza sample identified by MSDH occurred in week 12 (week ending March 28<sup>th</sup>).


The influenza cases were identified from the following counties: Alcorn (1), Attala (4), Benton (5), Bolivar (3), Choctaw (1), Coahoma (1), Copiah (3), Covington (2), DeSoto (2), Forrest (1), Franklin (3), Harrison (14), Hinds (3), Holmes (2), Humphreys (1), Itawamba (3), Jackson (1), Jefferson (2), Jefferson Davis (1), Jones (5), Kemper (3), Lafayette (3), Lauderdale (6), Leake (2), Lee (3), Leflore (1), Lowndes (5), Madison (9), Marion (5), Marshall (10), Monroe (5), Neshoba (5), Noxubee (3), Oktibbeha (3), Pearl River (5), Pike (2), Pontotoc (1), Prentiss (5), Rankin (9), Simpson (8), Sunflower (2), Tallahatchie (3), Tate (4), Tishomingo (2), Union (1), Walthall (1), Warren (2), Washington (7), Webster (9), Wilkinson (3), and Yazoo (6). Two cases were from unidentified counties. | Figure 5



#### **National and Mississippi Pediatric Mortality Surveillance**

Nationally, **no** influenza-associated pediatric deaths were reported to CDC during week **14**. To date, **125** influenza-associated pediatric deaths have been reported nationally during the 2014-2015 influenza season. | Figure 6






Of the **125** influenzaassociated pediatric deaths reported nationally during the 2014-2015 season, 103 (82%) have been attributed to influenza A viruses, 19 (15%) to influenza B viruses, two (2%) to an influenza A/B virus not distinguished, and one (1%) to an

influenza A and B virus co-infection. | Figure 7

Mississippi has had **one** influenza-associated pediatric death reported during this influenza season. The death occurred during week 10 (week ending March 14<sup>th</sup>). For additional information on influenza-associated pediatric deaths, please refer to the <u>CDC's FluView</u>.

#### **National ILI Surveillance**


For week **14**, the MS ILI rate (3.1%) remained **above** the national ILI rate (1.8%). | Figure 8



Mississippi reported "Sporadic" flu activity for week 14. | Table 3

Level of Flu Activity	Definition	
No Activity	Overall clinical activity remains low and there are no lab confirmed cases.	
Sporadic	Isolated cases of lab confirmed influenza in the state; ILI activity is not increased <u>OR</u> A lab-confirmed outbreak in a single institution in the state; ILI activity is not increased.	
Local	Increased ILI within a single region <b>AND</b> recent (within the past 3 weeks) laboratory evidence of influenza in that region. ILI activity in other regions is not increased <u>OR</u> two of more institutional outbreaks (ILI or lab confirmed) within a single region <b>AND</b> recent (within the past 3 weeks) lab confirmed influenza in that region. Other regions do not have increased ILI and virus activity is no greater than sporadic in those regions	
Regional	Increased ILI in at least 2 regions but fewer than half of the regions <b>AND</b> recent (within the past 3 weeks) lab confirmed influenza in the affected regions <u>OR</u> Institutional outbreaks (ILI or lab confirmed) in at least 2 regions but fewer than half of the regions <b>AND</b> recent lab confirmed influenza in the affected regions.	
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions <b>AND</b> recent (within the past 3 weeks) lab confirmed influenza in the state.	

During week 14, influenza activity continued to decrease in the United States. | Figure 9



<sup>&</sup>lt;sup>1</sup>For up-to-date information on flu activity nationwide, please refer to the CDC's website: <a href="http://www.cdc.gov/flu/weekly/fluactivitysurv.htm">http://www.cdc.gov/flu/weekly/fluactivitysurv.htm</a>.

#### **Additional information:**

Centers for Disease Control and Prevention	http://cdc.gov/flu/
Centers for Disease Control and Prevention FluView	http://www.cdc.gov/flu/weekly/
Flu.gov	http://www.flu.gov/
MSDH Flu and Pneumonia	http://msdh.ms.gov/msdhsite/_static/14,0,199.html
Google Flu Trends	http://www.google.org/flutrends/
World Health Organization FluNet	http://www.who.int/influenza/gisrs_laboratory/flunet/en/

Page intentionally blank

# **Appendix**

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9

